



A Physiologically Based Model of the HPG Axis in Fathead Minnows

Karen Watanabe, Zhenhong Li, Kevin Kroll, Natalia Reyero, Nancy Szabo,
Marisol Sepúlveda, Edward Orlando, Daniel Villeneuve, Michael Breen,
Gerald Ankley, and Nancy Denslow

Overview

- Background
 - The endocrine system
 - Endocrine active compounds (EACs) and their sources in the environment
- Objective
- Model Development
- Results
- Conclusions

The Endocrine System

- A network of glands distributed throughout the body

Hypothalamus

Pituitary

Pineal gland

Thyroid


Parathyroid

Adrenal

Pancreas

Gonad: Ovary or Testis

- These glands produce hormones that are distributed to distant target sites via the blood.



What are EACs?

- Chemicals, both natural and man-made, that can interfere with endocrine glands and their hormones or where the hormones act, i.e., the target tissues
- Environmental sources
 - Sewage treatment plant effluents
 - Pulp and paper mill effluents
 - Marine paints – tributyltin
 - Fungicides – vinclozolin and linuron
 - Herbicides – chlorotriazines such as atrazine
 - Plastics manufacturing - phthalates
 - Livestock industry pharmaceuticals

Effects in Wildlife

- Adverse effects observed in certain fish and wildlife
 - Estrogen exposure biomarker, vitellogenin, found in fish living near outflows from sewage treatment plants or pulp and paper mills
 - Imposex in molluscs exposed to tributyltin
 - Reproductive and immunological effects in marine mammals (PCBs and DDE)
 - Egg shell thinning and embryonic abnormalities in birds (DDT and PCBs)

Risk Assessment Needs

- Methods to predict effects of EACs on populations
- Tools to extrapolate effects from high to low dose
 - Target tissue dosimetry
 - Developing quantitative relationships between gene expression and higher order responses
- Identification of biomarkers as indicators of exposure and predictors of adverse effects
- Tools to extrapolate across species
 - Similarities across vertebrates for certain processes, e.g., steroidogenesis
 - Differences expected in ligand-receptor binding

Objective

- Develop a physiologically based computational model to simulate reproductive endpoints in male fathead minnows (FHM, *Pimephales promelas*) exposed to 17α -ethynylestradiol (EE_2)

Reproductive Endpoints of Interest

- Steroid hormones
 - Testosterone (T)
 - 11-ketotestosterone (KT)
 - 17 β -estradiol (E₂)
- Vitellogenin (Vtg)
 - Pre-cursor to a major egg yolk protein
 - Normally found at low levels in male FHM
 - Biomarker of exposure to estrogenic EACs because Vtg is induced after exposure


Why Model Fathead Minnows?

- Distributed throughout North America
 - Fresh water habitat
- Sexually dimorphic
- Fractional spawners
 - Spawn every 3-5 days
- Mature in 4-6 months
- EPA model for toxicity




Photo courtesy U.S. EPA

Why EE₂?





E₂
An estrogen synthesized
naturally



EE₂
Synthetic estrogen used in oral
contraceptives

Conceptual Model of the HPG Axis



Adapted from Villeneuve et al. (2007). Environmental Science & Technology 41(1): 321-330.

Physiologically Based Model Formulation

- Mass balance

F = volumetric flow


C = concentration

V = tissue compartment volume

R = rate of consumption (e.g., metabolism or binding), or a rate of production preceded by a “+” sign

t = time

$$\text{Accumulation} = \Sigma \text{ Flow in} - \Sigma \text{ Flow out} + \Sigma \text{ Production} - \Sigma \text{ Consumption}$$



$$\frac{d(C_{cv}V)}{dt} = F_{in} C_{in} - F_{out} C_{out} - R_c$$

Brain

$$\frac{d(V_{\text{brn}} C_{\text{EE2,brn}})}{dt} = F_{\text{brn}} (C_{\text{Art}_{\text{EE2}}} - C_{\text{Ven}_{\text{EE2,brn}}})$$


$$\begin{aligned} & -V_{\text{brn}} (k_{1\text{EE2,brn}} C_{\text{EE2,brn}} C_{\text{ERf,brn}} \\ & -k_{2\text{EE2:ER,brn}} C_{\text{EE2:ER,brn}}) \end{aligned}$$

$$\frac{d(V_{\text{brn}} C_{\text{E2,brn}})}{dt} = F_{\text{brn}} (C_{\text{Art}_{\text{E2}}} - C_{\text{Ven}_{\text{E2,brn}}})$$

$$\begin{aligned} & -V_{\text{brn}} (k_{1\text{E2,brn}} C_{\text{E2,brn}} C_{\text{ERf,brn}} \\ & -k_{2\text{E2:ER,brn}} C_{\text{E2:ER,brn}}) \end{aligned}$$

$$\begin{aligned} \frac{d(V_{\text{brn}} C_{\text{LH,brn}})}{dt} = & F_{\text{brn}} (C_{\text{Art}_{\text{LH}}} - C_{\text{Ven}_{\text{LH,brn}}}) \\ & + \rho_{\text{LH,brn}} C_{\text{E2,brn}} V_{\text{brn}} + \dot{m}_{\text{LH,brn}} \end{aligned}$$

Similar equations are formulated for T, KT, and Vtg



Gonad

$$\frac{d(V_{\text{gon}} C_{\text{EE2,gon}})}{dt} = F_{\text{gon}} (C_{\text{ArtEE2}} - C_{\text{VenEE2,gon}}) - V_{\text{gon}} (k_{1\text{EE2,gon}} C_{\text{EE2,gon}} C_{\text{ERf,gon}} - k_{2\text{EE2:ER,gon}} C_{\text{EE2:ER,gon}})$$

$$\begin{aligned} & - V_{\text{gon}} (k_{1\text{EE2,gon}} C_{\text{EE2,gon}} C_{\text{ERf,gon}} \\ & - k_{2\text{EE2:ER,gon}} C_{\text{EE2:ER,gon}}) \end{aligned}$$


$$\frac{d(V_{\text{gon}} C_{\text{E2,gon}})}{dt} = F_{\text{gon}} (C_{\text{ArtE2}} - C_{\text{VenE2,gon}}) - V_{\text{gon}} (k_{1\text{E2,gon}} C_{\text{E2,gon}} C_{\text{ERf,gon}} - k_{2\text{E2:ER,gon}} C_{\text{E2:ER,gon}})$$

$$\begin{aligned} & - V_{\text{gon}} (k_{1\text{E2,gon}} C_{\text{E2,gon}} C_{\text{ERf,gon}} \\ & - k_{2\text{E2:ER,gon}} C_{\text{E2:ER,gon}}) \\ & + V_{\text{max_arom}} C_{\text{T,gon}} / (K_{\text{m_arom}} + C_{\text{T,gon}}) \end{aligned}$$

$$\frac{d(V_{\text{gon}} C_{\text{LH,gon}})}{dt} = F_{\text{gon}} (C_{\text{ArtLH}} - C_{\text{VenLH,gon}}) - V_{\text{gon}} (k_{1\text{LH,gon}} C_{\text{LH,gon}} C_{\text{LR,gon}} - k_{2\text{LH:LR,gon}} C_{\text{LH:LR,gon}})$$

$$\begin{aligned} & - V_{\text{gon}} (k_{1\text{LH,gon}} C_{\text{LH,gon}} C_{\text{LR,gon}} \\ & - k_{2\text{LH:LR,gon}} C_{\text{LH:LR,gon}}) \end{aligned}$$

Similar equations are formulated for T, KT, and Vtg



Model Calibration Data


- Measurements for 154 unexposed male FHM (Dr. Ankley's Lab, U.S. EPA)
 - Body weight
 - Gonad weight
 - Liver weight
 - Reproductive endpoints include plasma concentrations of
 - E_2
 - T
 - KT
 - Vtg
- EE_2 48 hour exposure experiment
 - 50 ng/L in water
 - Measurements of
 - EE_2 concentration in pooled carcass and liver
 - Plasma T and Vtg concentrations

Model Input Parameter Estimation

- Rationale for parameter estimation by Markov Chain Monte Carlo (MCMC) simulation
 - Incorporates prior knowledge about the input parameters
 - Accounts for covariance between model input parameters
 - Results in a distribution of model parameters that can be used to quantify uncertainty in model predictions
- Computational requirements
 - Software: MCSim is available freely through GNU
 - Operating system: LINUX or UNIX



Markov Chain Monte Carlo

- 15 parameters estimated
 - Prior distributions informed by the literature
- Data from EE₂ exposed and unexposed FHM
 - Measured model outputs, y_j
 - E₂
 - T
 - Vitellogenin
 - EE₂



From Prior to Posterior Distributions

Prior distribution + Data \Rightarrow Posterior Distribution




Model Calibration


- Data
 - Half of the unexposed FHM data were used
 - All of the EE₂ experimental measurements were used
- MCMC simulation
 - 15 model parameters were calibrated
 - Four Markov chains were simulated containing 2500 iterations each
 - 1000 iterations required ~24 hours
 - Last 500 iterations from each chain were saved
 - Convergence evaluated using the potential scale reduction metric (Gelman et al., 1995)

Visualizing Convergence

Chains converge
Potential scale reduction = 1.0





Chains do not converge
Potential scale reduction = 4.2







Results: Parameter Estimation

EE₂ Partition Coefficient in FHM Gill (Blood to Water)



Results: Model Verification



Summary

- There is good agreement between median model predictions of plasma E_2 , KT, and Vtg concentrations
- The model underpredicts median plasma T concentrations
- The variance in model predictions for all four reproductive endpoints is lower than what is observed

Acknowledgments

- This research was supported in part by a Cooperative Agreement to the University of Florida, FL (ND), Oregon Health & Science University, OR (KW), Purdue University, IN (MS) and Florida Atlantic University, FL (EO) from the US Environmental Protection Agency (EPA) Science to Achieve Results program (STAR; Assistance Agreement # RD-83184801-0).
- Although the research described in this presentation has been funded wholly or in part by the U.S. Environmental Protection Agency, it has not been subjected to any formal EPA review and therefore does not necessarily reflect the views of the Agency, and no official endorsement should be inferred.
- ZL was supported by a grant from the Medical Research Foundation of Oregon.